

Short Communication

Selective Inhibition of Cytochrome P450 2D6 by Sarpogrelate and Its Active Metabolite, M-1, in Human Liver Microsomes[□]

Received August 18, 2013; accepted October 28, 2013

ABSTRACT

The present study was performed to evaluate the in vitro inhibitory potential of sarpogrelate and its active metabolite, M-1, on the activities of nine human cytochrome (CYP) isoforms. Using a cocktail assay, the effects of sarpogrelate on nine CYP isoforms and M-1 were measured by specific marker reactions in human liver microsomes. Sarpogrelate potently and selectively inhibited CYP2D6-mediated dextromethorphan *O*-demethylation with an IC₅₀ (K_i) value of 3.05 μM (1.24 μM), in a competitive manner. M-1 also markedly inhibited CYP2D6 activity; its inhibitory effect with an IC₅₀ (K_i) value of 0.201 μM (0.120 μM) was more potent than that of sarpogrelate, and was similarly potent as quinidine (K_i, 0.129 μM), a well-known typical CYP2D6 inhibitor. In addition, sarpogrelate and M-1 strongly inhibited both CYP2D6-catalyzed bufuralol 1'-hydroxylation and metoprolol

α-hydroxylation activities. However, sarpogrelate and M-1 showed no apparent inhibition of the other following eight CYPs: CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2E1, or CYP3A4/5. Upon 30-minute preincubation of human liver microsomes with sarpogrelate or M-1 in the presence of NADPH, no obvious shift in IC₅₀ was observed in terms of inhibition of the nine CYP activities, suggesting that sarpogrelate and M-1 are not time-dependent inactivators. Sarpogrelate strongly inhibited the activity of CYP2D6 at clinically relevant concentrations in human liver microsomes. These observations suggest that sarpogrelate could have an effect on the metabolic clearance of drugs possessing CYP2D6-catalyzed metabolism as a major clearance pathway, thereby eliciting pharmacokinetic drug-drug interactions.

Introduction

Sarpogrelate ((*R,S*)-1-[2-[2-(3-methoxyphenyl)ethyl]phenoxy]-3-(dimethyl amino)-2-propyl hydrogen succinate hydrochloride; Fig. 1) is a highly specific 5-HT_{2A} receptor antagonist widely used in China, Japan, and South Korea to treat peripheral arterial disease (Rashid et al., 2003; Doggrel, 2004). Sarpogrelate has inhibitory effects on serotonin-induced platelet aggregation (Hara et al., 1991a; Nakamura et al., 1999), thrombus formation (Hara et al., 1991b; Yamashita et al., 2000), vasoconstriction, and vascular smooth muscle cell proliferation (Sharma et al., 1999), all of which are mediated by 5-HT_{2A} receptors, and consequently reduces the ischemic symptoms associated with peripheral arterial disease. Additionally, sarpogrelate has beneficial effects in restenosis after coronary stenting (Fujita et al., 2003; Saini et al., 2004), pulmonary hypertension (Saini et al., 2004), angina pectoris (Kinugawa et al., 2002), and diabetes mellitus (Pietraszek et al., 1993; Ogawa et al., 1999), although the precise mechanisms remain unknown. Sarpogrelate is metabolized to (±)-1-[2-[2-(3-methoxyphenyl)ethyl]-phenoxy]-3-(dimethylamino)-2-propanol hydrochloride (M-1; Fig. 1), formed by hydrolysis from sarpogrelate (Nagatomo et al., 2004; Saini et al., 2004). The M-1 is an active sarpogrelate metabolite, which has inhibitory effects exceeding those of sarpogrelate in vitro (Pertz and Elz, 1995).

General sarpogrelate dosing in patients is one 100-mg tablet taken three times per day after meals (Shinohara et al., 2008). After oral administration of 100 mg of sarpogrelate to healthy male subjects, sarpogrelate is rapidly absorbed from the gastrointestinal tract with a mean maximum plasma concentration (C_{max}) of 856.3 ng/mL at 0.7 hour and is rapidly eliminated from plasma with a half-life of 0.8 hour (Kim et al., 2013a). The active metabolite, M-1, reaches a C_{max} of 49.3 ng/mL at 0.9 hour and exhibits slower elimination than sarpogrelate, with a half-life of 4.4 hours (Kim et al., 2013a). After absorption, sarpogrelate and M-1 further undergo glucuronide conjugations to form several metabolites, which are mainly excreted in bile (Kim et al., 2013b). Despite the wide use and excellent pharmacological properties of sarpogrelate, to date there is no information regarding the potential inhibitory effects of sarpogrelate and M-1 on human P450 isozymes.

In the present study, the inhibitory effects of sarpogrelate and M-1 on the nine cytochrome P450 (CYP) isozymes were evaluated using a cocktail assay to assess the potential of sarpogrelate to cause drug-drug interactions with other concomitantly administered drugs. We report in this work that especially M-1 is a selective competitive inhibitor of CYP2D6 in vitro.

Materials and Methods

Chemicals and Reagents. Pooled human liver microsomes from a mixed pool of 24 donors (17 males and 7 females), *S*-benzylmirtazapine, and 1'-hydroxybufuralol were purchased from BD Gentest (Woburn, MA). Sarpogrelate and M-1 were obtained from Kunwha Pharmaceutical Company (Seoul, Republic of Korea). Acetaminophen, bufuralol, chlorpropamide, chlorzoxazone, coumarin, dextropropofol, diethylthiocarbamate, furafylline, α-hydroxymetoprolol, ketoconazole, metoprolol, phenacetin, propranolol, quercetin, quinidine, rosiglitazone,

This work was supported by the Ministry of Trade, Industry, and Energy R&D Program, Republic of Korea [Grant 10039320], and 2013 Research Fund of Catholic University of Korea.

D.-Y.C. and S.H.B. contributed equally to this work.

dx.doi.org/10.1124/dmd.113.054296.

□ This article has supplemental materials available at dmd.aspetjournals.org.

ABBREVIATIONS: AUC, area under the plasma concentration-time curve; AUC_i, AUC in presence of an inhibitor; C_{max}, maximum plasma concentration; CYP, cytochrome P450; KF, potassium fluoride; LC-MS/MS, liquid chromatography-tandem mass spectrometry.




Fig. 1. Chemical structures of sarpogrelate (A) and its active metabolite, M-1 (B).

S-mephenytoin, sulfaphenazole, tolbutamine, 1,1',1''-phosphinothiolydinetriza-ziridine, potassium fluoride (KF), NADP, glucose 6-phosphate, glucose 6-phosphate dehydrogenase, and $MgCl_2$ were purchased from Sigma-Aldrich (St. Louis, MO). Bupropion, dextromethorphan, 6-hydroxy bupropion, 6-hydroxy-chlorzoxazone, 7-hydroxycoumarin, 4'-hydroxymephenytoin, 1'-hydroxymidazolam, *p*-hydroxy rosiglitazone, 4-hydroxytolbutamide, and midazolam were obtained from Toronto Research Chemicals (North York, ON, Canada). Solvents were high-performance liquid chromatographic grade (Burdick and Jackson Company, Morristown, NJ), and other chemicals were of the highest quality available.

Screening of Reversible Inhibitory Effects of Sarpogrelate and M-1 on the Activities of Nine Cytochrome P450 Enzymes. So-called cocktail assays in which several enzyme activities are determined in parallel by liquid chromatography-tandem mass spectrometry (LC-MS/MS) are particularly useful. The inhibitory potencies of sarpogrelate and M-1 were determined as described previously with slight modification (Bae et al., 2013). Phenacetin *O*-deethylase, coumarin 7-hydroxylase, bupropion 6-hydroxylase, rosiglitazone *p*-hydroxylase, tolbutamide 4-hydroxylase, 5-mephenytoin 4-hydroxylase, dextromethorphan *O*-demethylase, chlorzoxazone 6-hydroxylase, and midazolam 1'-hydroxylase activities were determined as probe activities in human liver microsomes for CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4/5, respectively. Sarpogrelate or M-1 (concentration: 0–50 μM) and all substrates were dissolved in acetonitrile and serially diluted with acetonitrile to the required concentrations to give a final organic solvent concentration of 1.0% in the incubation mixture. Concentrations of P450-selective substrates were used close to their reported K_m values (Supplemental Table 1; Kim et al., 2005; Bae et al., 2013).

Briefly, the incubation mixtures containing pooled human liver microsomes (final concentrations: 0.25 mg/ml), each P450-selective substrate, and a NADPH-generating system (1.3 mM $NADP^+$, 3.3 mM glucose 6-phosphate, 3.3 mM $MgCl_2$, and 0.4 U/ml glucose-6-phosphate dehydrogenase) were

preincubated for 5 minutes at 37°C. The reaction was initiated by adding an aliquot of sarpogrelate or M-1 (concentration range: 0–50 μM) and incubated for 15 minutes at 37°C in a shaking water bath. When sarpogrelate as an inhibitor was incubated, a 10- μL aliquot of 1 M KF in 0.1 M phosphate buffer (pH 7.4) was added before incubation to inhibit esterase activity (Clarke and Waskell, 2003). After incubation, reactions were stopped by addition of 50 μL of ice-cold acetonitrile containing 2 μM chlorpropamide as an internal standard, and they were chilled and centrifuged (13,000 rpm, 8 minutes, 4°C). The supernatant was then diluted 100-fold with acetonitrile and then injected into the LC-MS/MS system. All incubations were performed in triplicate, and mean values were used for analysis. Additionally, identical parallel incubation samples containing well-known reversible CYP inhibitors were included as positive controls. Two different microsomal protein concentrations, 0.05 and 0.1 mg/mL, were also used to evaluate the inhibitory potential for CYP2D6 activities.

Additionally, sarpogrelate or M-1 was tested as an inhibitor of bufuralol 1'-hydroxylase (Boobis et al., 1985; Kronbach et al., 1987), metoprolol α -hydroxylase (Otton et al., 1988), and other CYP2D6-specific biotransformation pathways. Concentrations of bufuralol (5 μM) and metoprolol (20 μM) were used in this study. Other procedures were similar to those used in the cocktail assays.

Determination of the K_i of Sarpogrelate and M-1 for CYP2D6. Based on the IC_{50} values, the K_i values of sarpogrelate and M-1 for CYP2D6 were determined. Briefly, dextromethorphan, a specific substrate for CYP2D6, was incubated with sarpogrelate, M-1, or quinidine, a well-known typical CYP2D6 inhibitor. For determination of K_i values, dextromethorphan concentrations used were 2.5, 5, and 10 μM . The concentrations of quinidine, sarpogrelate, and M-1 were as follows; 0–1 μM for quinidine, and 0–10 μM for sarpogrelate and M-1. All incubations were performed in triplicate, and mean values were used for the analysis. Other procedures were similar to those of the reversible inhibition studies.

Time-Dependent Inhibitory Effects of Sarpogrelate and M-1 on the Activities of Nine Cytochrome P450 Enzymes. The IC_{50} shift assay is one of the most efficient and convenient methods of evaluating the time-dependent inhibitory effects of sarpogrelate and M-1. Changes in enzymatic activity are usually detected with and without preincubation of the test compound for a defined period. A shift in IC_{50} to a lower value (shift) following preincubation indicates time-dependent inactivation (Obach et al., 2007).

Pooled human liver microsomes (1 mg/mL) were incubated with sarpogrelate or M-1 (0–50 μM) in the absence or presence of a NADPH-generating system for 30 minutes at 37°C (i.e., the inactivation incubation). After inactivation incubation, aliquots (10 μL) were transferred to fresh incubation tubes (final volume 100 μL) containing a NADPH-generating system and each P450-selective substrate cocktail set. When sarpogrelate was studied, a 10- μL aliquot of 1 M KF was added into both inactivation and incubation mixtures. The reaction system (100 μL total volume) was incubated for 15 minutes at 37°C in a shaking water bath. After incubation, reactions were stopped by addition of 50 μL of ice-cold acetonitrile containing 2 μM chlorpropamide, as an internal standard, and they were chilled and centrifuged (13,000 rpm, 8 minutes, 4°C). The supernatant was then diluted 10-fold with acetonitrile and injected into the LC-MS/MS system.

Determination of the Unbound Fraction of Sarpogrelate or M-1 in Human Liver Microsomes and Human Plasma. Equilibrium dialysis was conducted to assess the unbound fraction of sarpogrelate or M-1 in human liver microsomes and human plasma using a single-use plate rapid equilibrium dialysis device with dialysis membranes with a molecular weight cutoff of ~8000 Da (Thermo Scientific, Rockford, IL) (Bae et al., 2013). Human liver microsome samples containing sarpogrelate or M-1 at concentrations of 0.5 and 10 μM , respectively (100 μL), were dialyzed against 50 mM phosphate buffer (300 μL) at pH 7.4. The loaded dialysis plate was covered with sealing tape, placed on an orbital shaker at approximately 500 rpm, and incubated at 37°C for 4 hours. All incubations were performed in triplicate, and mean values were used for the analysis. Nonspecific binding in microsome/buffer mixed matrix was evaluated for sarpogrelate or M-1 concentrations using the LC-MS/MS method. In plasma protein-binding studies, the final concentrations of sarpogrelate or M-1 were both 0.5 and 10 μM . The LC-MS/MS conditions for determination of sarpogrelate and M-1 were optimized based on the conditions used in a previous study (Kim et al., 2013a).

LC-MS/MS Analysis. Metabolites of nine P450-selective substrates were analyzed using a tandem quadrupole mass spectrometer (QTrap 5500 LC-MS/MS; Applied Biosystems, Foster City, CA) equipped with an electrospray

ionization interface, as reported previously (Bae et al., 2013). Single reaction monitoring mode using specific precursor/product ion transition was used for quantification. The mass transitions of the metabolites of the nine P450-selective substrates and collision energies are listed in Supplemental Table 1. Peak areas for all of the analytes were integrated automatically using the Analyst software (version 1.5.2; Applied Biosystems, Foster City, CA).

The mass transitions used for quantification of 1'-hydroxybupifuralol or α -hydroxymetoprolol were optimized based on the conditions used in a previous study (VandenBrink et al., 2012).

Data Analysis. For reversible inhibition and time-dependent inhibition screening, the P450-mediated activities in the presence of the inhibitor, sarpogrelate or M-1, were expressed as percentages of the corresponding control values at 0 μ M sarpogrelate or M-1. From plots of percent inhibition versus inhibitor concentrations, corresponding IC_{50} values were calculated by nonlinear regression using the WinNonlin software (version 4.0; Pharsight, Mountain View, CA). The apparent kinetic parameters for inhibitory potential (K_i values) were estimated from the fitted curves using the WinNonlin software. The inhibition data were fit to different models of enzyme inhibition (competitive, noncompetitive, uncompetitive, or mixed) by nonlinear least-squares regression analysis (WinNonlin software). The most appropriate inhibition model selected based upon the goodness-of-fit criteria of a visual inspection of the data, correlation of determination (R^2), and corrected Akaike's Information Criterion.

For visual inspection, data are presented as Dixon plots and Lineweaver–Burk plots.

Results

The inhibitory effects of sarpogrelate and M-1 on the activities of nine CYP isozymes (CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4/5) at microsomal protein concentrations of 0.25 mg/mL are shown in Figs. 2 and 3, respectively. The IC_{50} values of sarpogrelate and M-1 at a microsomal protein concentration of 0.25 mg/mL are listed in Table 1. The IC_{50} values for the positive controls used in the reversible inhibition studies were in good agreement with published values to an acceptable degree of accuracy (data not shown). Of the nine P450 isoforms tested, CYP2D6-catalyzed dextromethorphan hydroxylation was most strongly inhibited by sarpogrelate and M-1, with IC_{50} values of 3.05 and 0.201 μ M, respectively (Table 1). Similar inhibitory potencies of sarpogrelate and M-1 at the different microsomal protein concentrations (0.05 and 0.1 mg/mL) were observed (data not shown), indicating that nonspecific microsomal binding of sarpogrelate and M-1 did not affect the inhibitory potencies. However, sarpogrelate and M-1 showed no apparent inhibition of the other eight




Fig. 2. IC_{50} curves of sarpogrelate for human P450 activities using the cocktail substrate, including CYP1A2 for phenacetin *O*-deethylase (A), CYP2A6 for coumarin 7-hydroxylase (B), CYP2B6 for bupropion hydroxylase (C), CYP2C8 for rosiglitazone *p*-hydroxylase (D), CYP2C9 for tolbutamide 4-hydroxylase (E), CYP2C19 for *S*-mephenytoin 4-hydroxylase (F), CYP2D6 for dextromethorphan *O*-demethylase (G), CYP2E1 for chlorzoxazone 6-hydroxylase (H), and CYP3A4/5 for midazolam 1'-hydroxylase (I). Data are the mean \pm SD of triplicate determinations. The dashed lines represent the best fit to the data using nonlinear regression.




Fig. 3. IC₅₀ curves of M-1 for human P450 activities using the cocktail substrate, including CYP1A2 for phenacetin *O*-deethylase (A), CYP2A6 for coumarin 7-hydroxylase (B), CYP2B6 for bupropion hydroxylase (C), CYP2C8 for rosiglitazone *p*-hydroxylase (D), CYP2C9 for tolbutamide 4-hydroxylase (E), CYP2C19 for *S*-mephenytoin 4-hydroxylase (F), CYP2D6 for dextromethorphan *O*-demethylase (G), CYP2E1 for chlorzoxazone 6-hydroxylase (H), and CYP3A4/5 for midazolam 1'-hydroxylase (I). Data are the mean \pm SD of triplicate determinations. The dashed lines represent the best fit to the data using nonlinear regression.

CYPs tested (Table 1; Figs. 2 and 3); the remaining activities at the tested highest concentration (50 μ M) were greater than 90%.

To determine whether the inhibitory activities of sarpogrelate and M-1 were substrate specific, we examined the inhibitory effects on other CYP2D6-specific biotransformation pathways (i.e., bufuralol 1'-hydroxylation and metoprolol α -hydroxylation) and found that sarpogrelate also markedly inhibited their activities, with IC₅₀ values of 4.02 and 3.37 μ M, respectively (data not shown). M-1 also potently inhibited CYP2D6 activity; corresponding IC₅₀ values were 0.360 and 0.545 μ M, respectively (data not shown).

To characterize the type of reversible inhibition of CYP2D6 by sarpogrelate or M-1 based on the IC₅₀ values, enzyme kinetic assays were conducted with varying concentrations of sarpogrelate or M-1 and dextromethorphan. Identical parallel incubation samples containing a known potent inhibitor of CYP2D6, quinidine, were included. The K_i values of sarpogrelate, M-1, and quinidine are listed in Table 2. Representative Dixon plots for the inhibition of CYP2D6 by sarpogrelate, M-1, and quinidine in human liver microsomes are shown in Fig. 4. Sarpogrelate and M-1 strongly and selectively inhibited CYP2D6 with K_i values of 1.24 μ M and 0.120 μ M, respectively. Specifically, inhibition of

CYP2D6 by M-1 was more potent than that of sarpogrelate and was similarly potent as quinidine (K_i , 0.129 μ M) (Table 3). Visual inspection of the Dixon plots and further analysis of the enzyme inhibition modes

TABLE 1

Values of IC₅₀ (μ M) of sarpogrelate and M-1 for each CYP isozyme in human liver microsomes

The assay conditions are described in *Materials and Methods*. Data are expressed as the mean of triplicate determinations.

CYPs	Reactions	IC ₅₀	
		Sarpogrelate	M-1
		μ M	
CYP1A2	Phenacetin <i>O</i> -deethylation	>50	>50
CYP2A6	Coumarin 7-hydroxylation	>50	>50
CYP2B6	Bupropion hydroxylation	>50	>50
CYP2C8	Paclitaxel 6 α -hydroxylation	>50	>50
CYP2C9	Tolbutamide 4-hydroxylation	>50	>50
CYP2C19	<i>S</i> -Mephenytoin 4'-hydroxylation	>50	>50
CYP2D6	Dextromethorphan <i>O</i> -demethylation	3.05	0.201
CYP2E1	Chlorzoxazone 6-hydroxylation	>50	>50
CYP3A4/5	Midazolam 1'-hydroxylation	>50	>50

TABLE 2

K_i values of the inhibition for CYP2D6 by sarpogrelate, M-1, and quinidine at microsomal protein concentrations of 0.25 mg/mL

Concentrations of sarpogrelate, M-1, and quinidine were as follows: 0–10 μM for sarpogrelate and M-1, and 0–1 μM for quinidine, respectively.

CYPs	K_i		
	Sarpogrelate	M-1	Quinidine
CYP2D6	1.24 ^a	0.120 ^a	0.129 ^a

^aInhibition type was determined by the best fit to competitive mode based on AICs.

suggested that the inhibition data of sarpogrelate, M-1, and quinidine all fit well to a competitive inhibition type.

A shift in the inhibition curve to a lower IC_{50} value by 30-minute preincubation in the presence of NADPH is an indicator of time-dependent inhibition. After 30-minute preincubation of sarpogrelate or M-1 with human liver microsomes in the presence of NADPH, no obvious shift in IC_{50} was observed for inhibition of the nine CYPs (data not shown). Representative IC_{50} shift plots for CYP2D6 activity by sarpogrelate or M-1 are shown in Supplemental Fig. 1. These suggest that sarpogrelate and M-1 are not time-dependent inhibitors.

The free fractions of sarpogrelate at concentrations of 0.5 and 10 μM in human plasma were $96.8 \pm 4.29\%$ and $95.1 \pm 3.12\%$, respectively ($n = 3$, each). However, incubation in human liver microsomes for 4 hours prohibited the measurement of microsomal binding for sarpogrelate due to its instability in microsomes. When M-1 was added at concentrations of 0.5 and 10 μM to human liver microsomes (human plasma), the free fractions of M-1 were $72.0 \pm 6.12\%$ ($97.3 \pm 4.28\%$) and $68.1 \pm 5.18\%$ ($97.8 \pm 3.09\%$), respectively ($n = 3$ each). The free fractions of sarpogrelate and M-1 were not affected by the concentrations added.

Discussion

To our knowledge, there are no reports of in vitro drug interactions of sarpogrelate via CYP isozymes. In this study, we demonstrated that sarpogrelate is a potent and selective competitive inhibitor of CYP2D6

in vitro. Additionally, M-1, an active metabolite of sarpogrelate, significantly inhibited CYP2D6 activities; its inhibitory effects with an IC_{50} (K_i) value of 0.201 μM (0.120 μM) were more potent than those of sarpogrelate, with an IC_{50} (K_i) value of 3.05 μM (1.24 μM). Sarpogrelate and M-1 strongly inhibited other CYP2D6-catalyzed bufuralol 1'-hydroxylation and metoprolol α -hydroxylation activities. However, sarpogrelate and M-1 showed no apparent inhibition of the other following eight CYPs: CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, or CYP2E1. Preincubation of sarpogrelate or M-1 with human liver microsomes and a NADPH-generating system did not alter the inhibition potencies against the nine CYPs, suggesting that sarpogrelate or M-1 are not time-dependent inactivators.

Generally, alterations in the activities of hepatic CYPs, inhibition or induction, in vivo represent the major mechanisms underlying pharmacokinetic drug–drug interactions (Clarke and Jones, 2002; Leucuta and Vlase, 2006). Although it accounts for only 2–5% of all hepatic CYP isozymes, CYP2D6 metabolizes approximately 25% of all clinically used medications, such as some cytotoxins, tamoxifen, and many agents used to treat associated complications such as antiarrhythmics, antiemetics, antidepressants, antipsychotics, and analgesics (Wolf and Smith, 1999; Ingelman-Sundberg and Evans, 2001). In addition, the CYP2D6 gene is highly polymorphic with more than 112 variants described to date (<http://www.imm.ki.se/CYPalleles/cyp2d6.htm>), and such variations in CYP2D6 expression are thought to increase the potential for drug–drug interactions (Bernard et al., 2006).

The in vitro inhibition potency alone does not dictate the likelihood of pharmacokinetic drug interactions because the in vivo concentration of the inhibitor should also be considered. For reversible inhibitors, the magnitude of the increase in exposure is related to the inhibitory potency (K_i), the concentration of inhibitor, and the fraction of the affected drug that ordinarily goes through the inhibited enzyme (f_m) (Yao & Levy, 2002; Ito et al., 2004; Obach et al., 2006). As stated previously, the C_{max} of sarpogrelate was 856.3 ng/mL (1.99 μM) following a single 100-mg sarpogrelate oral dose in healthy subjects. In addition, our clinical trial data indicate that the C_{max} values of sarpogrelate and M-1 in steady state are 657 ± 302 ng/ml (1.56 μM) at 0.9 hour with a half-life of 0.64 hour and 53.0 ± 16.1 ng/ml (0.161 μM)




Fig. 4. Dixon plots to determine K_i values for CYP2D6 of sarpogrelate (A), M-1 (B), and quinidine (C). The concentrations of dextromethorphan were determined 2.5 (●), 5 (○), and 10 (▼) μM , respectively. V represents formation rate of dextrophan (pmol/min/mg protein). Data are the mean values of triplicate determinations. The solid lines of sarpogrelate, M-1, and quinidine fit well to all competitive inhibition types.

TABLE 3

Prediction of maximum interactions with CYP2D6-cleared drugs caused by sarpogrelate

Values used for sarpogrelate include the following: $D = 100$ mg; $k_a = 0.0167/\text{min}$; $f_u = 0.04$, $C_{\text{max}} = 1.56$ μM (our unpublished data); $Q_h = 1470$ ml/min (Obach et al., 2006); adjusted K_i (as free fraction in microsomes) = 0.372 μM . The free fraction of sarpogrelate in microsomes is assumed to be same as that of M-1. The fraction of the affected drug cleared by CYP2D6 and F_a is assumed to be unity. The estimate for k_a was calculated from the expression $T_{\text{max}} = [\ln(k_a/k_c)]/(k_a - k_c)$, where k_c is the elimination rate constant. The value of k_c was calculated from half-life, 0.64 h, and T_{max} value was 0.9 h (our unpublished data).

$[I]_{\text{in vivo}}$	Equation Used to Estimate the $[I]_{\text{in vivo}}$	Fold Increase in Exposure (AUC _i /AUC) $1 + ([I]_{\text{in vivo}}/K_i)$
Systemic total C_{max}	$[I]_{\text{in vivo}} = C_{\text{max}}$ at steady state	5.19
Systemic free C_{max}	$[I]_{\text{in vivo}} = f_u \times C_{\text{max}}$ at steady state	1.17
Estimated total portal C_{max}	$[I]_{\text{in vivo}} = C_{\text{max}} + k_a \times F_a \times D/Q_h$	11.5
Estimated total free C_{max}	$[I]_{\text{in vivo}} = f_u \times (C_{\text{max}} + k_a \times F_a \times D/Q_h)$	1.42

D, dose of inhibitor; F_a , fraction absorbed from the gastrointestinal tract; f_u , unbound fraction of sarpogrelate in plasma; $[I]$, concentrations of sum of inhibitors; k_a , absorption rate constant; Q_h , human hepatic blood flow rate.

at 1.08 hours with a half-life of 4.98 hours, respectively (our unpublished data).

In human in vivo interaction studies, the degree of interaction is usually expressed as the ratio of the area under the plasma concentration–time curve (AUC) in the presence (AUC_i) and absence of an inhibitor. When the in vivo inhibition potency of sarpogrelate against completely CYP2D6-cleared drug ($f_m = 1$) is determined from the plasma concentration of sarpogrelate described above, the K_i values of sarpogrelate (1.24 μM) for CYP2D6, and the unbound fractions in both human liver microsomes and plasma by the methods of Obach et al. (2006), the AUC_i to AUC ratio is estimated to be 1.17–11.5 (Table 3). These estimates of the magnitude of drug–drug interactions for a CYP2D6-cleared drug range from 1.17 to 11.5, largely due to whether the unbound or total sarpogrelate concentrations are most relevant to enzyme inhibition in vivo. However, all the AUC_i to AUC ratios had >1.1 when either total or unbound concentrations were used for the calculation of ratios. Thus, we cannot exclude the possibility of the in vivo inhibitory potency of CYP2D6 by sarpogrelate.

There are some limitations to our calculations. First, the effects of M-1 were not considered, although its inhibitory effect was more potent than that of sarpogrelate. Second, the free fraction of sarpogrelate in human liver microsomes is assumed to be the same as that of M-1. Instability of sarpogrelate in microsomes for 4-hour incubation did not allow for an accurate measurement. Finally, from a clinical drug–drug interaction point of view, a meaningful inhibitory effect of sarpogrelate might not be observed despite high inhibitory potencies of sarpogrelate (K_i 1.24 μM) and M-1 (K_i 0.120 μM) for CYP2D6 due to their considerable short half-lives (sarpogrelate, 0.64 hour and M-1, 4.98 hours) and T_{max} (sarpogrelate, 0.9 hour and M-1, 1.08 hour). This may explain previously observed interactions between paroxetine as a potent CYP2D6 inhibitor and clozapine as a CYP2D6 substrate (Hiemke and Härter, 2000). Applying dosages above 20 mg paroxetine per day produced a substantial increase in clozapine plasma levels (Centorrino et al., 1996), whereas a fixed dose of 20 mg/d could not observe increased plasma levels of clozapine (Wetzel et al., 1998). The magnitude of CYP2D6 inhibition correlates with its plasma concentrations (Hiemke and Härter, 2000). Therefore, further investigations are required to clarify the in vivo interactions of CYP2D6-targeted drugs with sarpogrelate.

In conclusion, these observations suggest that sarpogrelate and M-1 are potent and selective competitive inhibitors of CYP2D6 in vitro. Especially, inhibition of CYP2D6 by M-1 was 10-fold more potent than that of sarpogrelate, and was similarly potent as quinidine, a well-known typical CYP2D6 inhibitor. The data support the use of M-1 as a well-known inhibitor of CYP2D6 instead of quinidine for routine screening of P450 reversible inhibition when human liver

microsomes are used as the enzyme source. Finally, it would be expected that sarpogrelate could have an effect on the metabolic clearance of drugs possessing CYP2D6-catalyzed metabolism as a major clearance pathway, thereby eliciting pharmacokinetic drug–drug interactions.

Department of Family Practice and Community Health, Ajou University School of Medicine, Suwon, Republic of Korea (D.-Y.C., B.-T.K.); College of Pharmacy and Integrated Research Institute of Pharmaceutical Sciences, Catholic University of Korea, Bucheon, Republic of Korea (S.H.B., J.K.L., S.K.B.); and Department of Emergency Medicine, Inje University College of Medicine, Busan, Republic of Korea (Y.W.K.)

DOO-YEOUN CHO
SOO HYEON BAE
JOENG KEE LEE
YANG WEON KIM
BOM-TAECK KIM
SOO KYUNG BAE

Authorship Contributions

Participated in research design: Cho, Y.W. Kim, B.-T. Kim, S.H. Bae, S.K. Bae.

Conducted experiments: S.H. Bae, Lee.

Contributed analytic tools: S.H. Bae, Lee.

Performed data analysis: Cho, S.H. Bae, Y.W. Kim, B.-T. Kim, S.K. Bae.

Wrote or contributed to the writing of the manuscript: Cho, S.H. Bae, Y.W. Kim, B.-T. Kim, S.K. Bae.

References


- Bae SH, Kwon MJ, Choi EJ, Zheng YF, Yoon KD, Liu KH, and Bae SK (2013) Potent inhibition of cytochrome P450 2B6 by sibutramine in human liver microsomes. *Chem Biol Interact* **205**: 11–19.
- Bernard S, Neville KA, Nguyen AT, and Flockhart DA (2006) Interethnic differences in genetic polymorphisms of CYP2D6 in the U.S. population: clinical implications. *Oncologist* **11**: 126–135.
- Boobis AR, Murray S, Hampden CE, and Davies DS (1985) Genetic polymorphism in drug oxidation: in vitro studies of human debrisoquine 4-hydroxylase and bufuralol 1'-hydroxylase activities. *Biochem Pharmacol* **34**:65–71.
- Centorrino F, Baldessarini RJ, Frankenburg FR, Kando J, Volpicelli SA, and Flood JG (1996) Serum levels of clozapine and norclozapine in patients treated with selective serotonin reuptake inhibitors. *Am J Psychiatry* **153**:820–822.
- Clarke SE and Jones BC (2002) Human cytochromes P450 and their role in metabolism-based drug–drug interactions, in *Drug-Drug Interactions* (Rodrigues AD ed) pp 55–88, Marcel Dekker, New York.
- Clarke TA and Waskell LA (2003) The metabolism of clopidogrel is catalyzed by human cytochrome P450 3A and is inhibited by atorvastatin. *Drug Metab Dispos* **31**:53–59.
- Doggrell SA (2004) Sarpogrelate: cardiovascular and renal clinical potential. *Expert Opin Investig Drugs* **13**:865–874.
- Fujita M, Mizuno K, Ho M, Tsukahara R, Miyamoto A, Miki O, Ishii K, and Miwa K (2003) Sarpogrelate treatment reduces restenosis after coronary stenting. *Am Heart J* **145**:E16.
- Hara H, Kitajima A, Shimada H, and Tamao Y (1991a) Antithrombotic effect of MCI-9042, a new antiplatelet agent on experimental thrombosis models. *Thromb Haemost* **66**:484–488.
- Hara H, Osakabe M, Kitajima A, Tamao Y, and Kikumoto R (1991b) MCI-9042, a new antiplatelet agent is a selective 52-serotonergic receptor antagonist. *Thromb Haemost* **65**:415–420.

- Hiemke C and Härter S (2000) Pharmacokinetics of selective serotonin reuptake inhibitors. *Pharmacol Ther* **85**:11–28.
- Ingelman-Sundberg M and Evans WE (2001) Unravelling the functional genomics of the human CYP2D6 gene locus. *Pharmacogenetics* **11**:553–554.
- Ito K, Brown HS, and Houston JB (2004) Database analyses for the prediction of in vivo drug-drug interactions from in vitro data. *Br J Clin Pharmacol* **57**:473–486.
- Kim HJ, Jeong ES, Seo KA, Shin KJ, Choi YJ, Lee SJ, Ghim JL, Sohn DR, Shin JG, and Kim DH (2013b) Glucuronidation of a sarpogrelate active metabolite is mediated by UDP-glucuronosyltransferases 1A4, 1A9, and 2B4. *Drug Metab Dispos* **41**:1529–1537.
- Kim MJ, Kim H, Cha IJ, Park JS, Shon JH, Liu KH, and Shin JG (2005) High-throughput screening of inhibitory potential of nine cytochrome P450 enzymes in vitro using liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom* **19**:2651–2658.
- Kim TE, Kim JR, Jung JA, Kim SR, Lee JW, Jun H, Lee SY, Huh W, and Ko J (2013a) Comparison of pharmacokinetics between sarpogrelate hydrochloride immediate-release formulation and controlled-release formulation. *Int J Clin Pharmacol Ther* **51**:114–119.
- Kinugawa T, Fujita M, Lee JD, Nakajima H, Hanada H, and Miyamoto S (2002) Effectiveness of a novel serotonin blocker, sarpogrelate, for patients with angina pectoris. *Am Heart J* **144**:E1.
- Kronbach T, Mathys D, Gut J, Catin T, and Meyer UA (1987) High-performance liquid chromatographic assays for bufuralol 1'-hydroxylase, debrisoquine 4-hydroxylase, and dextromethorphan O-demethylase in microsomes and purified cytochrome P-450 isozymes of human liver. *Anal Biochem* **162**:24–32.
- Leucuta SE and Vlase L (2006) Pharmacokinetics and metabolic drug interactions. *Curr Clin Pharmacol* **1**:5–20.
- Nagatomo T, Rashid M, Abul Muntasir H, and Komiya T (2004) Functions of 5-HT_{2A} receptor and its antagonists in the cardiovascular system. *Pharmacol Ther* **104**:59–81.
- Nakamura K, Kariyazono H, Moriyama Y, Toyohira H, Kubo H, Yotsumoto G, Taira A, and Yamada K (1999) Effects of sarpogrelate hydrochloride on platelet aggregation, and its relation to the release of serotonin and P-selectin. *Blood Coagul Fibrinolysis* **10**:513–519.
- Obach RS, Walsky RL, and Venkatakrishnan K (2007) Mechanism-based inactivation of human cytochrome p450 enzymes and the prediction of drug-drug interactions. *Drug Metab Dispos* **35**:246–255.
- Obach RS, Walsky RL, Venkatakrishnan K, Gaman EA, Houston JB, and Tremaine LM (2006) The utility of in vitro cytochrome P450 inhibition data in the prediction of drug-drug interactions. *J Pharmacol Exp Ther* **316**:336–348.
- Ogawa S, Takeuchi K, Sugimura K, Sato C, Fukuda M, Lee R, Ito S, and Sato T (1999) The 5-HT₂ receptor antagonist sarpogrelate reduces urinary and plasma levels of thromboxane A₂ and urinary albumin excretion in non-insulin-dependent diabetes mellitus patients. *Clin Exp Pharmacol Physiol* **26**:461–464.
- Otton SV, Crewe HK, Lennard MS, Tucker GT, and Woods HF (1988) Use of quinidine inhibition to define the role of the sparteine/debrisoquine cytochrome P450 in metoprolol oxidation by human liver microsomes. *J Pharmacol Exp Ther* **247**:242–247.
- Pertz H and Elz S (1995) In-vitro pharmacology of sarpogrelate and the enantiomers of its major metabolite: 5-HT_{2A} receptor specificity, stereoselectivity and modulation of ritanserin-induced depression of 5-HT contractions in rat tail artery. *J Pharm Pharmacol* **47**:310–316.
- Pietraszek MH, Takada Y, Taminato A, Yoshimi T, Watanabe I, and Takada A (1993) The effect of MCI-9042 on serotonin-induced platelet aggregation in type 2 diabetes mellitus. *Thromb Res* **70**:131–138.
- Rashid M, Manivet P, Nishio H, Pratuangdejkul J, Rajab M, Ishiguro M, Launay JM, and Nagatomo T (2003) Identification of the binding sites and selectivity of sarpogrelate, a novel 5-HT₂ antagonist, to human 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C} receptor subtypes by molecular modeling. *Life Sci* **73**:193–207.
- Saini HK, Takeda N, Goyal RK, Kumamoto H, Arneja AS, and Dhalla NS (2004) Therapeutic potentials of sarpogrelate in cardiovascular disease. *Cardiovasc Drug Rev* **22**:27–54.
- Sharma SK, Zahradka P, Chapman D, Kumamoto H, Takeda N, and Dhalla NS (1999) Inhibition of serotonin-induced vascular smooth muscle cell proliferation by sarpogrelate. *J Pharmacol Exp Ther* **290**:1475–1481.
- Shinohara Y, Nishimaru K, Sawada T, Terashi A, Handa S, Hirai S, Hayashi K, Tohgi H, Fukuuchi Y, and Uchiyama S, et al.; S-ACCESS Study Group (2008) Sarpogrelate-aspirin comparative clinical study for efficacy and safety in secondary prevention of cerebral infarction (S-ACCESS): a randomized, double-blind, aspirin-controlled trial. *Stroke* **39**:1827–1833.
- VandenBrink BM, Foti RS, Rock DA, Wienkers LC, and Wahlstrom JL (2012) Prediction of CYP2D6 drug interactions from in vitro data: evidence for substrate-dependent inhibition. *Drug Metab Dispos* **40**:47–53.
- Wetzel H, Anghelescu I, Szegedi A, Wiesner J, Weigmann H, Härter S, and Hiemke C (1998) Pharmacokinetic interactions of clozapine with selective serotonin reuptake inhibitors: differential effects of fluvoxamine and paroxetine in a prospective study. *J Clin Psychopharmacol* **18**:2–9.
- Wolf CR and Smith G (1999) Cytochrome P450 CYP2D6. *IARC Sci Publ* **148**:209–229.
- Yamashita T, Kitamori K, Hashimoto M, Watanabe S, Giddings JC, and Yamamoto J (2000) Conjoint effects of the 5HT₂ receptor antagonist, sarpogrelate, on thrombolysis with modified tissue plasminogen activator in different laser-induced thrombosis models. *Haemostasis* **30**:321–332.
- Yao C and Levy RH (2002) Inhibition-based metabolic drug-drug interactions: predictions from in vitro data. *J Pharm Sci* **91**:1923–1935.

Address correspondence to: Dr. Soo Kyung Bae, College of Pharmacy and Integrated Research Institute of Pharmaceutical Sciences, Catholic University of Korea, 43 Jibong-ro, Wonmi-gu, Bucheon 420-743, Republic of Korea. E-mail: baesk@catholic.ac.kr

Selective inhibition of cytochrome P450 2D6 by sarpogrelate and its active metabolite, M-1, in human liver microsomes. Doo-Yeoun Cho, Soo Hyeon Bae, Joeng Kee Lee, Yang Weon Kim, Bom-Taeck Kim, Soo Kyung Bae. *Drug Metabolism and Disposition*

Supplemental Figure 1.



Supplemental Figure 1. Representative IC_{50} shift plots for CYP2D6 by sarpogrelate (A) and M-1 (B) with human liver microsomes in the presence (\bullet , the dashed lines) and absence (\circ , the solid lines) of NADPH for 30 min pre-incubation. Data are the mean \pm SD of triplicate determinations. The dashed lines and solid lines represent the best fit to the data using non-linear regression.

Selective inhibition of cytochrome P450 2D6 by sarpogrelate and its active metabolite, M-1, in human liver microsomes. Doo-Yeoun Cho, Soo Hyeon Bae, Joeng Kee Lee, Yang Weon Kim, Bom-Taeck Kim, Soo Kyung Bae. *Drug Metabolism and Disposition*

Supplemental Table 1. Optimized mass parameters for the detection of metabolites of the nine P450-selective substrates and internal standard using cocktail assays on the API5500 Qtrap instruments

P450s	Substrates	Concentration (μM)	Metabolites	Transitions (m/z)	ESI mode	DP (eV)	CE (eV)
1A2	Phenacetin	50	Acetaminophen	152 / 110	+	50	20
2A6	Coumarin	5	7-Hydroxycoumarin	161 / 105	-	-90	-30
2B6	Bupropion	50	6-Hydroxybupropion	256 / 238	+	140	20
2C8	Rosiglitazone	1	<i>p</i> -Hydroxyrosiglitazone	374 / 151	+	60	30
2C9	Tolbutamide	100	4-Hydroxytolbutamide	285 / 104	-	-20	-40
2C19	<i>S</i> -Mephenytoin	100	4'-Hydroxymepheytoin	233 / 190	-	-45	-20
2D6	Dextromethorphan	5	Dextrophan	258 / 157	+	190	50
2E1	Chlorzoxazone	50	6-Hydroxychlorzoxazone	184 / 120	-	-100	-40
3A	Midazolam	5	1'-Hydroxymidazolam	342 / 203	+	20	40
Internal standard	Chlorpropamide			277 / 111	+	120	20
				275 / 190	-	-80	-50

ESI, electrospray ionization; DP, declustering potential; CE, collision energy.

The Optimized ion spray voltage and temperature were set at ± 5500 V and 500°C , respectively.